arXiv:gr-qc/0501077v1 26 Jan 2005

Exact solutions of the Dirac equation in Robertson-Walker space-time

Xin-Bing Huang*

Department of Physics, Peking University,

100871, Beijing, P.R.China

Abstract

The covariant Dirac equation in Robertson-Walker space-time is studied under the comoving coordinates. The exact forms of the spatial factor of wave function are respectively acquired in closed, spatially flat, and open universes.

PACS numbers: 03.65.Ge, 04.20.Jb, 03.65.Pm, 98.80.Jk

---

*huangxb@pku.edu.cn

1 Introduction

The Robertson-Walker metric plays a central role in modern cosmology. The expanding universe is usually described by it. Therefore the behavior of relativistic particles obeying the covariant Dirac equation in Robertson-Walker space-time has evoked many investigations in history. Such works go back to Schrödinger [1], Brill and Wheeler [2], Barut and Duru [3]. A general discussion of the case of massless neutrino has been given by Brill and Wheeler.

In literature, Li and Guo ever considered the exact solutions of Klein-Gordon equation and the Dirac equation in Beltrami-de-Sitter coordinates [4]. They gave right solutions for Klein-Gordon equation and acquired the discrete spectrum [5] of the mass of a spin-0 particle because of the continuous condition of the Legend function at the boundary of de Sitter space-time. But they obtained the wrong evolutionary equation from the Dirac equation, thus gave the wrong solutions for the Dirac equation in de Sitter space-time.

Had been aware of the importance of the behavior of a spin-$\frac{1}{2}$ particle in the expanding universe, Barut and Duru investigated the exact solutions of the Dirac equation for three typical models of expanding universe [3]. But they assumed that the space-time is spatially flat. Thus the exact solution of the Dirac equation in general Robertson-Walker space-time haven't been given in literature.

In this paper, we investigate the behavior of a massive spin-$\frac{1}{2}$ particle in the open, spatially flat, and closed Robertson-Walker space-time, respectively. The exact forms of the spatial factor are obtained. Our work will be a correction and extension to the paper of Li and Guo, on the other hand it is also a complementary to the work of Barut and Duru. The exact solutions of the Dirac equation in Robertson-Walker will give a stage for studying the possible usage of neutrino oscillations [6] in cosmology, we think.

2 Dirac equation in Robertson-Walker space-time

The cosmological principle, the hypothesis that all positions in the universe are essentially equivalent, assumes that the "smeared" universe is homogeneous and isotropic about every point. The convenient implement to describe this "smeared" universe is the Robertson-Walker metric, which is manifestly spherically symmetric. In the comoving coordinates, the Robertson-Walker metric of the universe is taken to be[†]

$$ds^2 = \mathrm{g}_{\mu\nu} dx^\mu dx^\nu = dt^2 - R^2(t) \left[\frac{dr^2}{1-kr^2} + r^2(d\theta^2 + \sin^2\theta d\varphi^2) \right] , \tag{1}$$

where $R(t)$ is an unknown function of time, which is called cosmic scale factor, and k is a constant, which by a suitable choice of units for r can be chosen to have the value +1, 0, or −1. When $k = +1,\ 0$, or -1, the space-time is respectively closed, spatially flat or open.

To set the stage for solving the Dirac equation in a curved space-time, we introduce the local tetrad field as follows[‡]

$$\mathrm{g}^{\mu\nu} = e^\mu{}_a(x) e^\nu{}_b(x) \eta^{ab} , \tag{2}$$

where $\mathrm{g}^{\mu\nu}$ being the space-time metric, and the metric tensor of Minkowski space-time η_{ab} is

$$\eta^{00} = +1 , \qquad \eta^{11} = \eta^{22} = \eta^{33} = -1 , \qquad \eta^{ab} = 0 \quad \text{for} \quad a \neq b . \tag{3}$$

In a curved space-time, the covariant Dirac equation reads

$$\left[i\gamma^\mu(x) \frac{\partial}{\partial x^\mu} - i\gamma^\mu(x)\Gamma_\mu(x) \right] \psi(x) = m\psi(x) , \tag{4}$$

Here $\gamma^\mu(x)$ are the curvature-dependent Dirac matrices and $\Gamma_\mu(x)$ are the spin connections to be determined. The curvature-dependent Dirac matrices $\gamma^\mu(x)$ are presented in terms of the tetrad field as

$$\gamma^\mu(x) = e^\mu{}_a(x)\gamma^a , \tag{5}$$

[†] we adopt the conventions that $x^0 = t$, $x^1 = r$, $x^2 = \theta$, $x^3 = \varphi$ and $\hbar = c = 1$.

[‡] In this paper, using Roman suffixes to refer to the bases of local Minkowski frame; using Greek suffixes to refer to curvilinear coordinates of space-time.

where γ^a denotes the standard flat-space Dirac matrices, which satisfies $\{\gamma^a, \gamma^b\} = 2\eta^{ab}$.

The tetrad field in the case of Robertson-Walker metric (1) is thus taken to be

$$e^{\mu}{}_{a}(x) = \mathrm{diag}\left(1, \frac{\sqrt{1-kr^2}}{R(t)}, \frac{1}{R(t)r}, \frac{1}{R(t)r\sin\theta}\right) \ . \tag{6}$$

Inserting Eq.(6) into Eq.(5) yields

$$\begin{aligned} \gamma^0(x) &= \gamma_0 \ , \qquad \gamma^1(x) = -\frac{\sqrt{1-kr^2}}{R(t)}\gamma_1 \ , \\ \gamma^2(x) &= -\frac{1}{R(t)r}\gamma_2 \ , \quad \gamma^3(x) = -\frac{1}{R(t)r\sin\theta}\gamma_3 \ . \end{aligned} \tag{7}$$

The spin connections $\Gamma_\mu(x)$ in Eq.(4) satisfy the equation

$$[\Gamma_\mu(x), \gamma^\nu(x)] = \frac{\partial\gamma^\nu(x)}{\partial x^\mu} + \Gamma^\nu_{\mu\rho}\gamma^\rho(x) \ , \tag{8}$$

where $\Gamma^\nu_{\mu\rho}$ are the Christoffel symbols for the metric (1), which are determined by

$$\Gamma^\nu_{\mu\rho} = \frac{1}{2}\mathrm{g}^{\nu\tau}\left(\frac{\partial \mathrm{g}_{\tau\rho}}{\partial x^\mu} + \frac{\partial \mathrm{g}_{\mu\tau}}{\partial x^\rho} - \frac{\partial \mathrm{g}_{\mu\rho}}{\partial x^\tau}\right) \ . \tag{9}$$

Hence Eq.(8) can be solved for the spin connections $\Gamma_\mu(x)$ which we determine as

$$\begin{aligned} \Gamma_0 &= 0 \ , \qquad \Gamma_1 = \frac{\dot{R}(t)}{2\sqrt{1-kr^2}}\gamma_0\gamma_1 \ , \\ \Gamma_2 &= \frac{1}{2}\dot{R}(t)r\gamma_0\gamma_2 + \frac{1}{2}\sqrt{1-kr^2}\gamma_2\gamma_1 \ , \\ \Gamma_3 &= \frac{1}{2}\dot{R}(t)r\sin\theta\gamma_0\gamma_3 + \frac{1}{2}\sqrt{1-kr^2}\sin\theta\gamma_3\gamma_1 + \frac{1}{2}\cos\theta\gamma_3\gamma_2 \ , \end{aligned} \tag{10}$$

where $\dot{R}(t)$ denotes $\frac{dR(t)}{dt}$, so that the combination $\gamma^\mu(x)\Gamma_\mu(x)$ in Eq.(4) simplifies to

$$\gamma^\mu(x)\Gamma_\mu(x) = -\frac{3\dot{R}}{2R}\gamma_0 + \frac{\sqrt{1-kr^2}}{R(t)r}\gamma_1 + \frac{\cos\theta}{2R(t)r\sin\theta}\gamma_2 \ . \tag{11}$$

Inserting Eq.(7) and Eq.(11) into Eq.(4), then we obtain the Dirac equation in Robertson-Walker space-time

$$\begin{aligned} &\left\{iR\gamma_0\left(\frac{\partial}{\partial t} + \frac{3\dot{R}}{2R}\right) - i\left[\sqrt{1-kr^2}\gamma_1\left(\frac{\partial}{\partial r} + \frac{1}{r}\right) + \frac{1}{r}\gamma_2\left(\frac{\partial}{\partial\theta} + \frac{\cos\theta}{2\sin\theta}\right)\right.\right. \\ &\qquad\left.\left. + \frac{1}{r\sin\theta}\gamma_3\frac{\partial}{\partial\varphi}\right] - mR\right\}\psi(t,r,\theta,\varphi) = 0 \ . \end{aligned} \tag{12}$$

3 The exact solutions of the Dirac equation

We can simplify the Dirac equation (12) by writing [4]

$$\psi(t,r,\theta,\varphi) = (\sin\theta)^{-\frac{1}{2}} R^{-\frac{3}{2}} \Psi(t,r,\theta,\varphi) \ . \tag{13}$$

The reduced equation in terms of $\Psi(t,r,\theta,\varphi)$ is of the form

$$R\left(\frac{\partial}{\partial t} + im\gamma_0\right)\Psi = \left[\sqrt{1-kr^2}\gamma_0\gamma_1\left(\frac{\partial}{\partial r} + \frac{1}{r}\right) + \frac{1}{r}\gamma_1\hat{K}(\theta,\varphi)\right]\Psi \ , \tag{14}$$

where

$$\hat{K}(\theta,\varphi) = \gamma_0\gamma_1\left(\gamma_2\frac{\partial}{\partial\theta} + \gamma_3\frac{1}{\sin\theta}\frac{\partial}{\partial\varphi}\right) \tag{15}$$

is a Hermitian operator, as we know, which is defined in Ref.[7] first. The eigen equation of $\hat{K}$ is

$$\hat{K}\Psi_{\varsigma,\varepsilon} = \varsigma\Psi_{\varsigma,\varepsilon} \ , \qquad \varsigma = 0, \pm 1, \pm 2, \cdots, \tag{16}$$

where the eigen value ς is an integer, which has been proven in Ref.[2, 7].

Only the two matrices γ_0 and γ_1 remain explicitly in the simplified Dirac equation (14), they can therefore be represented by 2×2 matrices

$$\gamma_0 = \begin{pmatrix} \mathbf{1} & 0 \\ 0 & -\mathbf{1} \end{pmatrix} , \quad \gamma_1 = \begin{pmatrix} 0 & -\tau_1 \\ \tau_1 & 0 \end{pmatrix} , \tag{17}$$

where $\mathbf{1}$, τ_1 are respectively 2×2 unit matrix and the Pauli matrix, namely

$$\tau_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} . \tag{18}$$

We separate the angular factor from the wave function, and represent the radial and temporal factor by a two-component spinor, that is

$$\Psi_{\varsigma,\varepsilon} = \Theta_\varsigma(\theta,\varphi)\begin{pmatrix} \phi(r,t) \\ \chi(r,t) \end{pmatrix} , \tag{19}$$

where the angular factor, $\Theta_\varsigma(\theta,\varphi)$, is determined by the requirement $\hat{K}\Theta_\varsigma(\theta,\varphi) = \varsigma\Theta_\varsigma(\theta,\varphi)$. This equation for eigenstates of the angular motion has been investigated by Schrödinger [7]. He found, the operator $\hat{K}$ is related to the total angular momentum. Their eigenvalues satisfy [8]

$$\varsigma = \mp(j+\frac{1}{2}) = \begin{cases} -(l+1) & \text{for} \quad j = l + \dfrac{1}{2} \\ l & \text{for} \quad j = l - \dfrac{1}{2} \end{cases} . \tag{20}$$

Also the eigenfunction Θ_ς is related with the spherical harmonics as follows

$$\Theta_\varsigma(\theta,\varphi) \propto Y_{l,\bar{m}+\frac{1}{2}}(\theta,\varphi) \propto P_l^{\bar{m}+\frac{1}{2}}(\cos\theta) e^{i(\bar{m}+\frac{1}{2})\varphi} , \tag{21}$$

or

$$\Theta_\varsigma(\theta,\varphi) \propto Y_{l,\bar{m}-\frac{1}{2}}(\theta,\varphi) \propto P_l^{\bar{m}-\frac{1}{2}}(\cos\theta) e^{i(\bar{m}-\frac{1}{2})\varphi} , \tag{22}$$

where $P(\cos\theta)$ being the associated Legendre functions, and $\bar{m} \pm \frac{1}{2} = 0, \pm 1, \pm 2, \cdots, \pm l$.

Using equations (16), (17), and (19), we can rewrite the equation (14) into two equations

$$R\left(\frac{\partial}{\partial t} + im\right)\phi(r,t) = \left[\sqrt{1-kr^2}\left(\frac{\partial}{\partial r} + \frac{1}{r}\right) + \frac{\varsigma}{r}\right]\tau_1 \chi(r,t) , \tag{23}$$

$$R\left(\frac{\partial}{\partial t} - im\right)\chi(r,t) = \left[\sqrt{1-kr^2}\left(\frac{\partial}{\partial r} + \frac{1}{r}\right) - \frac{\varsigma}{r}\right]\tau_1 \phi(r,t) . \tag{24}$$

To solve the above equations, we separate the functions $\phi(t,r)$ and $\chi(t,r)$ into radial and temporal factors, respectively

$$\phi(t,r) = U_1(r)T_1(t) , \qquad \chi(t,r) = U_2(r)T_2(t). \tag{25}$$

With the help of Eq.(18), inserting the above equations into Eq.(23) and Eq.(24) yields the evolution equations

$$R\left(\frac{d}{dt} + im\right)T_1(t) - i\varepsilon T_2(t) = 0 , \tag{26}$$

$$R\left(\frac{d}{dt} - im\right)T_2(t) - i\varepsilon T_1(t) = 0 , \tag{27}$$

and the radial equations

$$\left[\sqrt{1-kr^2}\left(\frac{d}{dr} + \frac{1}{r}\right) + \frac{\varsigma}{r}\right]\left[\sqrt{1-kr^2}\left(\frac{d}{dr} + \frac{1}{r}\right) - \frac{\varsigma}{r}\right]U_1 + \varepsilon^2 U_1 = 0 , \tag{28}$$

$$\left[\sqrt{1-kr^2}\left(\frac{d}{dr} + \frac{1}{r}\right) - \frac{\varsigma}{r}\right]\left[\sqrt{1-kr^2}\left(\frac{d}{dr} + \frac{1}{r}\right) + \frac{\varsigma}{r}\right]U_2 + \varepsilon^2 U_2 = 0 . \tag{29}$$

Obviously, if U_1 is substituted by U_2, and simultaneously ς are replaced by $-\varsigma$, then the equation (28) becomes the equation (29). Therefore, in the following, it is of only necessity to study the equation (28).

The equation (28) directly simplifies to

$$(1-kr^2)\frac{d^2U_1}{dr^2}+\left(\frac{2}{r}-3kr\right)\frac{dU_1}{dr}+\left(\frac{\varsigma\sqrt{1-kr^2}-\varsigma^2}{r^2}+\varepsilon^2-k\right)U_1=0\ . \tag{30}$$

The constant k appears in the above equation, the value of which will affect the exact solution of Eq.(30). We, therefore, investigate Eq.(30) according to the value of k respectively.

3.1 In spatially flat Robertson-Walker geometry

In the case of spatially flat Robertson-Walker geometry, the constant k satisfies $k=0$. The equation (30) is thus reduced to

$$\frac{d^2U_1}{dr^2}+\frac{2}{r}\frac{dU_1}{dr}+\left(\frac{\varsigma(1-\varsigma)}{r^2}+\varepsilon^2\right)U=0\ . \tag{31}$$

Through a simple transformation, one can easily find that the equation (31) is the Bessel's equation of order $-\varsigma+\frac{1}{2}$. The solutions of the above equation are [9]

$$U_1(r)=\frac{1}{\sqrt{\varepsilon r}}J_{\pm(\varsigma-\frac{1}{2})}(\varepsilon r)\ . \tag{32}$$

Where $J_{\pm(\varsigma-\frac{1}{2})}(\varepsilon r)$ are called the Bessel functions.

3.2 In closed Robertson-Walker geometry

In the case of $k=+1$, the Robertson-Walker space-time is closed. The equation (30) becomes

$$(1-r^2)\frac{d^2U_1}{dr^2}+\left(\frac{2}{r}-3r\right)\frac{dU_1}{dr}+\left(\frac{\varsigma\sqrt{1-r^2}-\varsigma^2}{r^2}+\varepsilon^2-1\right)U_1=0\ . \tag{33}$$

Do the transformation

$$\rho=\frac{\sqrt{1-r^2}-1}{\sqrt{1-r^2}+1}\ , \tag{34}$$

then, in terms of ρ, the radial equation (33) is rewritten as follows

$$\rho(1-\rho)^2\frac{d^2U_1}{d\rho^2}+\frac{1}{2}(1-\rho)(3-\rho)\frac{dU_1}{d\rho}+\left[\frac{\varsigma(1-\rho)}{2\rho}-\frac{(\varsigma+\varsigma^2)(1-\rho)^2}{4\rho}+1-\varepsilon^2\right]U_1=0\ . \tag{35}$$

The above equation can be transformed into a hypergeometric equation, In the interval of $0\le r<1$, the solutions of which can thus be represented as [9]

$$\begin{aligned} U_1(r,\varsigma>0) &= \left(\frac{\sqrt{1-r^2}-1}{\sqrt{1-r^2}+1}\right)^{\frac{\varsigma-1}{2}}\left(\frac{2}{\sqrt{1-r^2}+1}\right)^{1+\varepsilon} \\ &\quad \cdot F\left(\varsigma+\varepsilon+\frac{1}{2},\varepsilon,\varsigma+\frac{1}{2};\frac{\sqrt{1-r^2}-1}{\sqrt{1-r^2}+1}\right)\ , \\ U_1(r,\varsigma\le 0) &= \left(\frac{\sqrt{1-r^2}-1}{\sqrt{1-r^2}+1}\right)^{-\frac{\varsigma}{2}}\left(\frac{2}{\sqrt{1-r^2}+1}\right)^{1+\varepsilon} \\ &\quad \cdot F\left(1+\varepsilon,\frac{1}{2}+\varepsilon-\varsigma,\frac{3}{2}-\varsigma;\frac{\sqrt{1-r^2}-1}{\sqrt{1-r^2}+1}\right)\ . \end{aligned} \tag{36}$$

Where F is the hypergeometric function.

3.3 In open Robertson-Walker geometry

In this subsection, we investigate the case of $k=-1$, where the space-time is open. The radial equation (30) is simplified to

$$(1+r^2)\frac{d^2U_1}{dr^2}+\left(\frac{2}{r}+3r\right)\frac{dU_1}{dr}+\left(\frac{\varsigma\sqrt{1+r^2}-\varsigma^2}{r^2}+\varepsilon^2+1\right)U_1=0\ . \tag{37}$$

Do the similar transformation as Eq.(34), set

$$\rho=\frac{\sqrt{1+r^2}-1}{\sqrt{1+r^2}+1}\ . \tag{38}$$

We then rewrite the equation (37) as

$$\rho(1-\rho)^2\frac{d^2U_1}{d\rho^2}+\frac{1}{2}(1-\rho)(3-\rho)\frac{dU_1}{d\rho}+\left[\frac{\varsigma(1-\rho)}{2\rho}-\frac{(\varsigma+\varsigma^2)(1-\rho)^2}{4\rho}+\varepsilon^2+1\right]U_1=0\ . \tag{39}$$

Similar to the equation (35), the above equation can also be transformed into a hypergeometric equation [9], the solutions of which are taken to be

$$U_1(r,\varsigma>0) = \left(\frac{\sqrt{1+r^2}-1}{\sqrt{1+r^2}+1}\right)^{\frac{\varsigma-1}{2}}\left(\frac{2}{\sqrt{1+r^2}+1}\right)^{1+i\varepsilon}$$

$$
\begin{aligned}
& \cdot F\left(\varsigma+i\varepsilon+\frac{1}{2}, i\varepsilon, \varsigma+\frac{1}{2}; \frac{\sqrt{1+r^2}-1}{\sqrt{1+r^2}+1}\right) \quad , \\
U_1(r,\varsigma\leq 0) \quad = \quad & \left(\frac{\sqrt{1+r^2}-1}{\sqrt{1+r^2}+1}\right)^{-\frac{\varsigma}{2}}\left(\frac{2}{\sqrt{1+r^2}+1}\right)^{1+i\varepsilon} \\
& \cdot F\left(1+i\varepsilon, \frac{1}{2}+i\varepsilon-\varsigma, \frac{3}{2}-\varsigma; \frac{\sqrt{1+r^2}-1}{\sqrt{1+r^2}+1}\right) \quad . \qquad (40)
\end{aligned}
$$

We have indicated that the solution of U_2 can be obtained from equations (32), (36), and (40) by substituting $-\varsigma$ for ς. Hence we have acquired the exact forms of the spatial factor of wave function in Robertson-Walker geometry.

In a different coordinates of spatially flat Robertson-Walker space-time, the authors of Ref.[3] have investigated the exact solutions of the Dirac equation for three models of expanding universes. It is obvious that our evolutionary equations (26) and (27) are equivalent to the equation (10) in Ref.[3]. Hence the solutions of equations (26) and (27) must be the same as the results given in Ref.[3].

The authors of Ref.[3] have given the exact solutions of three special models for expansion: 1. $R = a_0 t$, a_0 being a constant, a model of a linear expanding universe; 2. $R = a_0\sqrt{t}$, a model of a radiation-dominated universe; 3. $R = e^{Ht}$, H being a constant also, a model of de Sitter universe. These models are typical in cosmology, we will not discuss any more.

4 Conclusion

The behavior of the spin-$\frac{1}{2}$ particles in Robertson-Walker space-time is investigated. The exact expressions for the spatial factor of the spin-$\frac{1}{2}$ wave functions in closed, spatially flat, and open Robertson-Walker space-time are given, respectively. This work is a complementary to the Ref.[3].

Acknowledgement: I would like to thank Prof. D. Du, Prof. C. J. Zhu, Prof. Z. Chang, Prof. A. Sugamoto and Prof. I. Oda for their help and encouragement.

References

[1] Schrödinger E 1932 Preuss. Akad. Wiss. Berlin, Ber. **11-12** 105; 1939 Physica **6** 899; 1940 Proc. R. Ir. Acad., Sect. A **46** 9

[2] Brill D R and Wheeler J A 1957 Rev. Mod. Phys. **29** 465

[3] Barut A O and Duru I H 1987 Phys. Rev. D **36** 3705

[4] Li G Y and Guo H Y 1982 Acta Physica Sinica (in Chinese) **31** 1501

[5] Chang Z and Huang X B 2002 Phys. Letts. A **297** 320

[6] Xing Z Z 2004 Int. J. Mod. Phys. A **19** 1

[7] Schrödinger E 1938 Commentationes Pontif. Acad. Sci. **2** 321

[8] Greiner W 2000 *Relativistic quantum mechanics* (third edition, Springer, Hong Kong), pp.210-214

[9] Wang Z X and Guo D R 2000 *Introduction to Special Function* (Peking University Press, Beijing)